

# Formation of star-forming clouds from the magnetised, diffuse interstellar medium

Robi Banerjee & Bastian Körtgen

Hamburger Sternwarte, Universität Hamburg,  
Gojenbergsweg 112, 21029 Hamburg, Germany

Molecular clouds, the birthplaces of stars in galaxies, form dynamically from the diffuse atomic gas of the interstellar medium (ISM). The ISM is also threaded by magnetic fields which have a large impact on its dynamics. In particular, star forming regions must be magnetically *supercritical* in order to accommodate gas clumps which can collapse under their own weight. Based on a parameter study of three dimensional magneto-hydrodynamical (MHD) simulations, we show that the long-standing problem of how such supercritical regions are generated is still an open issue.

## 1 Introduction

Present day stars form within the densest regions of molecular clouds (MCs) and giant molecular clouds (GMCs), in gravitationally unstable cores and clumps. Our common understanding is that those MCs and GMCs form from the diffuse, atomic (HI) gas within timescales of less than 10 Myr<sup>1,2</sup>. The generation of filaments and substructures within GMCs is primarily controlled by magnetic fields and turbulence<sup>3,4,5</sup>. In particular, magnetic fields are an elemental part of the interstellar medium<sup>6,7,8,9,10,11</sup> which have a large impact on the dynamics of the ISM on various spatial scales<sup>12,13,14,15</sup> as the magnetic energy density is comparable to the thermal energy density of the ISM<sup>16</sup>.

One long-standing issue is the formation of supercritical clumps and cores. Similarly to thermal pressure, magnetic fields prevent contraction of otherwise (thermally) self-gravitating gas clumps if the magnetic fields are strong enough. Therefore, gaseous overdensities must be magnetically *supercritical*, quantified by the mass-to-flux ratio,  $\mu$ , to collapse and to subsequently allow the formation of stars.

Already in 1956, Mestel & Spitzer realised that molecular clouds should be magnetically *subcritical* assuming field strengths that correspond to the equipartition of magnetic and kinetic energy density within the ISM<sup>17</sup>. To generate supercritical cloud cores out of those subcritical conditions, they suggested that the non-perfect coupling between charged particles and neutrals, i.e. the ambipolar diffusion (AD) drift, could locally reduce the mass-to-flux ratio which allows the cloud to break up and to form stars. For a long time this was the standard theory of star formation out of the magnetised ISM<sup>3,18</sup>. In this fairly static “standard model” of magnetically-supported, AD-mediated supercritical cores, low-mass stars would form by the slow gravitational contraction of isolated cores containing a very small fraction of the clouds’ mass. This picture would also account for the very low observed global star formation efficiency (SFE) of giant molecular clouds<sup>19,20</sup>. The slow contraction results from the typical timescale for ambipolar diffusion,  $t_{AD}$ , which is an order of magnitude larger than the free-fall time,  $t_{ff}$ , of individual cloud cores (their ratio is about  $t_{AD}/t_{ff} \approx 10 (x_e/10^{-7})$ , where  $x_e$  is the ionisation fraction). On the other




Figure 1. These observational data summarise the main motivation for our proposed study: How do subcritical (HI) clouds become supercritical (H<sub>2</sub>) clouds? Our previous studies have shown that it is everything but trivial to build up supercritical clouds out of the magnetised interstellar medium, because the mass-to-flux ratio is fairly well conserved, even in the presence of ambipolar diffusion and enhanced non-ideal MHD setups<sup>45,46</sup>. [From Crutcher (2012)<sup>7</sup>]

hand, rather recent observations by Crutcher (2009)<sup>21</sup> of individual cloud cores including Zeeman measurements to determine their magnetic field distribution indicate that idealised models of ambipolar-diffusion driven star formation are unlikely to be operative. In idealised models with ordered background magnetic fields, efficient ambipolar diffusion would lead to a local increase of the mass-to-flux ratio towards the centre of cloud cores which is not seen in their observed sample<sup>21</sup> (but see also Bertram et al. (2012)<sup>22</sup> on the difficulty to interpret those observations).

However, present-day models of star formation also account for the fact that molecular clouds are also pervaded by supersonic random motions, i.e. turbulence<sup>23,24,25</sup>. Eventually, this resulted in a paradigm shift of the theory of star formation where magnetic fields only play a minor role and supersonic, super-Alfvénic turbulence controls the star formation efficiency within molecular clouds<sup>4,26,27,28,29</sup>. As a consequence, the magnetic fields are expected to be highly disordered rather than being an ordered background field. Hence, idealised models of ambipolar diffusion drift should not apply<sup>30,31</sup>. Additionally, the AD characteristic timescale is expected to decrease in this case<sup>32,33,34,35</sup> and other diffusive effects like turbulent reconnection might be operative<sup>36,37</sup>. Indeed, a number of studies have suggested that both MCs<sup>38</sup> and their clumps<sup>39,40,41,42,43,44</sup> are close to being magnetically critical, with a moderate preference for being supercritical. Moreover, recent compilations of observational data show that cloud cores and clumps with column densities of  $N \gtrsim 2 \times 10^{21} \text{ cm}^{-2}$  are essentially *all* supercritical (see Fig. 1).

Whether those supercritical cloud cores and clumps are the result of ambipolar diffusion together with random motions in the ISM is far from being certain and has to be investigated further. For instance, recently Heitsch & Hartmann (2014)<sup>47</sup> argued in their




Figure 2. Setup of the initial conditions. *Left*: The well studied head-on collision where both flows are in the direction of the background magnetic field. *Right*: Flows with an inclination angle with respect to the background magnetic field. This setup resembles the impact of external driving of such flows (e.g. driving by SN blast waves). [From Körtgen & Banerjee (2015)<sup>46</sup>]

parameter study that ambipolar diffusion in concert with turbulence is unlikely to control the formation of supercritical cores and hence star formation. They again propose an alternative scenario where large scale flows are the main driver to generate supercritical cores. This idea, where supercritical clouds could be assembled from large scale flows was already discussed in Mestel & Spitzer (1956)<sup>17</sup> as an alternative to the AD-mediated scenario and to avoid the “magnetic flux problem”. But only in combination with supersonic turbulence this scenario becomes more feasible because gravitational fragmentation could be suppressed during the assembly of the clouds by those turbulent motions<sup>48,49</sup>. This accumulation idea would also support a number of recent observations which show that magnetic fields are dynamically important on all scales in the Milky Way and other spiral galaxies<sup>50,12,14,51,13,15</sup>. This is particular evident from Fig. 1: The low column density HI gas is magnetically *subcritical*, whereas clouds which exceed columns of  $N \gtrsim 2 \times 10^{21} \text{ cm}^{-2}$  are magnetically *supercritical*.

In the presented numerical parameter study, we investigated the possibility of diffusion mediated generation of *supercritical* clouds showing that it is unlikely that such unstable clouds can be build up from *subcritical* HI-clouds.

## 2 Numerical method and initial conditions

For these studies we used the FLASH adaptive mesh refinement (AMR) code<sup>52,53</sup>. In addition to the basic ideal MHD equations (for which we employ the Bouchut solver<sup>54,55,56</sup>) we also used the *ambipolar diffusion* module developed from Duffin & Pudritz (2008)<sup>57</sup>. Additionally, self-gravity as well as heating and cooling processes were included in those simulations. For the latter, we followed the treatment by Koyama & Inutsuka (2002)<sup>58</sup> (an analytic simplification of their detailed calculation in<sup>59,60,61</sup>). To capture the build-up of self-gravitating cores within the molecular clouds we used *sink particles*<sup>62</sup> in addition to the Jeans refinement criterion (i.e. the Truelove criterion<sup>63</sup>). In particular the detailed

sink particle approach allows us to unambiguously identify supercritical, collapsing regions which are important for our studies quantifying the star formation ability from the magnetised ISM.

Our initial setups for those studies are similar to the ones described in<sup>60,61,45</sup> (see also Fig. 2) where the build-up of molecular clouds is modeled by the collision of cylindrical streams of warm neutral HI gas (WNM). Each flow is  $l = 112$  pc long and has a radius of  $r = 64$  pc. The bulk flows are slightly supersonic with typical Mach numbers of  $\mathcal{M}_f = 2$ . On top of those bulk motions, a turbulent velocity field is superimposed which triggers initial instabilities like the non-linear thin-shell instability (NTSI)<sup>64</sup> and subsequently leads to fragmentation of the cloud. The initially uniform magnetic field has a strength of  $B = \{3, 4, 5\}$   $\mu\text{G}$  corresponding to mass-to-flux ratios of  $\mu/\mu_{\text{crit}} \approx 1, 0.7, 0.6$  if the critical value  $\mu_{\text{crit}} \approx 0.13/\sqrt{G}$  is applied<sup>65</sup>.

Furthermore, we also studied the impact of an oblique angle of the flows with respect to the background magnetic field (see right panel of Fig. 2). Those oblique flows are more realistic than the head-on flows and could be generated, for instance, by supernova shock waves and by the gravitational potential of spiral arms. The motion of the flow at an inclination with respect to the magnetic field results in enhanced magnetic diffusivity (by numerical diffusion<sup>66,67,68</sup>). Again, those flows resemble streams of the WNM in a thermally bistable configuration<sup>69</sup>. The flows are studied with different oblique angles, which are varied from  $10^\circ$  to  $60^\circ$ , different initial magnetic fields strengths and different strength of the initial turbulence ranging from subsonic to supersonic velocity fluctuations. For details on the numerical setup and our initial conditions see Körtgen & Banerjee (2015)<sup>46</sup>.

### 3 Results

As can be seen from Fig. 3, the different initial field strengths have significant implications for the resulting dynamical behaviour of the molecular cloud. The main difference comes about in efficiency to form stars (or not). In the case of a rather weak background field of 3  $\mu\text{G}$  supercritical star forming clumps can be generated whereas in the case of a slightly stronger, but more realistic, magnetic field star formation is fully suppressed. Note that, due to the oblique flows with an angle of  $60^\circ$  the effective mass-to-flux ratios are 0.73 in the 3  $\mu\text{G}$  case and 0.44 in the 5  $\mu\text{G}$  case. That means that both cases are initially subcritical, but only in the cases of the weak magnetic field locally supercritical clumps are assembled due to sufficient flux loss.

An interesting point is also the field morphology. In the weak magnetic field case the field structure in the dense regions is clearly separated from the large scale magnetic field, whereas in the strong field case the field morphology is almost unaffected compared to the initial configuration (see the blue stream lines of Fig. 3). From an observational point of view, the field structure and its dynamical importance within molecular clouds is still debated. On the one hand, some multi-scale polarisation data indicate that magnetic fields in GMCs are essentially just dragged in from larger scales and are dynamically important<sup>70,12,13,51</sup>. On the other hand, Zeeman measurements of individual cloud cores together with analyses of numerical simulations indicate rather weak fields that might not be dynamically important<sup>21,71</sup>. With our subsequent studies on cloud formation on cloud scales including a more detailed modelling of ambipolar diffusion we hope to clarify this issue.




Figure 3. Results from colliding flow simulations investigating the formation of molecular clouds. Here, the flows collide with an oblique angle of  $60^\circ$ . *Left panel:* The weak field case ( $3 \mu\text{G}$ ). The *Right panel* shows the same situations in the case of a stronger background magnetic field of  $5 \mu\text{G}$ . In the case of a weak magnetic field supercritical cloud cores can form that allow the formation of stars (marked with black dots). Stronger initial magnetic fields prohibit the formation of stars even in the case of large oblique angles of the flows. The blue stream-lines indicate the magnetic field morphology in the projected 2D plane. [From Körtgen & Banerjee (2015)<sup>46</sup>]

In Fig. 4 we quantify the main results by means of histograms in the  $N$ - $B$ -plane from our colliding flow studies for various initial conditions. Only in the *initially* marginally subcritical case ( $B = 3 \mu\text{G}$ ) we observe signs of star formation within supercritical cores<sup>a</sup>. For slightly stronger initial magnetic fields ( $B \gtrsim 4 \mu\text{G}$ ) no supercritical cloud cores are generated, hence there is no star formation activity, regardless whether ambipolar diffusion is active or the flows collide with an oblique angle. Nevertheless the results of those simulation show the observed behaviour in the low column regime ( $N \lesssim 10^{21} \text{ cm}^{-2}$ ), where gas assembles along field lines without changing the field strength by much (see also the latest analysis from PLANCK observations of individual molecular clouds)<sup>51</sup>. Only within supercritical, self-gravitating cores the magnetic field gets enhanced by compression due to flux freezing. Furthermore, we observe that star formation is immediately initiated, when the gas becomes supercritical promoting a picture of “rapid” star formation<sup>49</sup>.

## 4 Conclusions

Here, we summarised our recent results from MHD simulations of colliding flows with varying initial conditions on the possible formation of supercritical cloud cores from subcritical initial conditions. Although, dense clouds are easily formed within colliding flow scenarios due to thermal instability, the generation of supercritical clumps are largely determined by the initial conditions. Furthermore, increasing initial turbulence lead to lower masses of the cores and clumps because the HI streams become less coherent. Otherwise, increasing magnetic field strength lead to more massive molecular clouds, which nevertheless do not become supercritical. Oblique flows still lead to cloud cores with masses

<sup>a</sup>If we assume  $\mu_{\text{crit}} \approx 0.13/\sqrt{G}$  for spherical cores<sup>65</sup> we get  $\mu/\mu_{\text{crit}} = 0.97 (3 \mu\text{G}/B)$  for our head-on colliding flow configurations.




Figure 4. Results from colliding flow simulations with various different initial conditions. Shown are histograms of the line-of-sight field strength  $B_{\text{LOS}}$  as function of the column density  $N$ . From left to right:  $B = 3 \mu\text{G}$ ,  $B = 4 \mu\text{G}$  and  $B = 5 \mu\text{G}$ , respectively. Top:  $\Phi = 0^\circ$ , Bottom:  $\Phi = 60^\circ$ . Different line colours denote different times. Also shown are the criticality condition (red line<sup>30,7</sup>), corrected for projection effects (black line<sup>72</sup>), and assuming equipartition of turbulent and magnetic fields (blue line<sup>73</sup>). Colour coded is the mass distribution within this two parameter space. [from Körtgen & Banerjee (2015)<sup>46</sup>]

comparable to what has been observed recently. But starting with subcritical HI flows, in no case the magnetic flux loss is sufficient to allow the build-up of supercritical cloud cores. Generally, increasing inclination of the flows lead to increasing diffusivity of the magnetic field. Again, regardless of the variation of the inclination, no tendency for faster accumulation of gas or faster transition to thermally dominated regions was seen in our simulations.

We therefore stress the role of magnetic fields in the context of molecular cloud and star formation. We point out the complete lack of supercritical regions for realistic initial field strengths. From the observational side, HI clouds may be supercritical as a whole, but their observed, dense subregions be subcritical.

Hence, the question remains, how magnetically *supercritical* cloud cores are formed?

## Acknowledgments

RB acknowledges funding by the DFG via the Emmy-Noether grant BA 3706/1-1, the ISM-SPP 1573 grants BA 3706/3-1 and BA 3706/3-2, as well as for the grant BA 3706/4-1. The software used in this work was in part developed by the DOE-supported ASC/Alliance Center for Astrophysical Thermonuclear Flashes at the University of Chicago. The authors gratefully acknowledge the Gauss Centre for Supercomputing (GCS) for providing computing time through the John von Neumann Institute for Computing (NIC) on the GCS share of the supercomputer JUQUEEN<sup>74</sup> at

Jülich Supercomputing Centre (JSC). GCS is the alliance of the three national supercomputing centres HLRS (Universitt Stuttgart), JSC (Forschungszentrum Jülich), and LRZ (Bayerische Akademie der Wissenschaften), funded by the German Federal Ministry of Education and Research (BMBF) and the German State Ministries for Research of Baden-Württemberg (MWK), Bayern (StMWFK) and Nordrhein-Westfalen (MIWF).

## References

1. Williams, J. P., Blitz, L. & McKee, C. F. The Structure and Evolution of Molecular Clouds: from Clumps to Cores to the IMF. *Protostars and Planets IV* 97 (2000). [astro-ph/9902246](#).
2. Blitz, L. *et al.* Giant Molecular Clouds in Local Group Galaxies. *Protostars and Planets V* 81–96 (2007). [arXiv:astro-ph/0602600](#).
3. Shu, F. H., Adams, F. C. & Lizano, S. Star formation in molecular clouds - Observation and theory. *ARA&A* **25**, 23–81 (1987).
4. Mac Low, M.-M. & Klessen, R. S. Control of star formation by supersonic turbulence. *Reviews of Modern Physics* **76**, 125–194 (2004).
5. André, P. *et al.* From Filamentary Networks to Dense Cores in Molecular Clouds: Toward a New Paradigm for Star Formation. *Protostars and Planets VI* 27–51 (2014). [1312.6232](#).
6. Beck, R. Magnetic Fields in Galaxies. *Space Sci. Rev.* **166**, 215–230 (2012).
7. Crutcher, R. M. Magnetic Fields in Molecular Clouds. *ARA&A* **50**, 29–63 (2012).
8. Heiles, C. & Haverkorn, M. Magnetic Fields in the Multiphase Interstellar Medium. *Space Sci. Rev.* **166**, 293–305 (2012).
9. Beck, R. & Wielebinski, R. *Magnetic Fields in Galaxies*, 641 (Springer Science+Business Media Dordrecht, 2013).
10. Li, H.-B. *et al.* The Link Between Magnetic Fields and Cloud/Star Formation. *Protostars and Planets VI* 101–123 (2014). [1404.2024](#).
11. Li, Z.-Y. *et al.* The Earliest Stages of Star and Planet Formation: Core Collapse, and the Formation of Disks and Outflows. *Protostars and Planets VI* 173–194 (2014). [1401.2219](#).
12. Li, H.-B. & Henning, T. The alignment of molecular cloud magnetic fields with the spiral arms in M33. *Nature* **479**, 499–501 (2011). [1111.2745](#).
13. Li, H.-B. *et al.* Self-similar fragmentation regulated by magnetic fields in a region forming massive stars. *Nature* **520**, 518–521 (2015).
14. Pillai, T. *et al.* Magnetic Fields in High-mass Infrared Dark Clouds. *ApJ* **799**, 74 (2015). [1410.7390](#).
15. Franco, G. A. P. & Alves, F. O. Tracing the magnetic field morphology of the Lupus I molecular cloud. *ArXiv 1504.08222*, *accepted for publication in ApJ* (2015). [1504.08222](#).
16. Heiles, C. & Troland, T. H. The Millennium Arecibo 21 Centimeter Absorption-Line Survey. IV. Statistics of Magnetic Field, Column Density, and Turbulence. *ApJ* **624**, 773–793 (2005). [arXiv:astro-ph/0501482](#).
17. Mestel, L. & Spitzer, L., Jr. Star formation in magnetic dust clouds. *MNRAS* **116**, 503 (1956).
18. Mouschovias, T. C. & Ciolek, G. E. Magnetic Fields and Star Formation: A Theory

- Reaching Adulthood. In C. J. Lada & N. D. Kylafis (ed.) *NATO ASIC Proc. 540: The Origin of Stars and Planetary Systems*, 305–+ (1999).
19. Myers, P. C. *et al.* Molecular clouds and star formation in the inner galaxy - A comparison of CO, H II, and far-infrared surveys. *ApJ* **301**, 398–422 (1986).
  20. Evans, N. J., II *et al.* The Spitzer c2d Legacy Results: Star-Formation Rates and Efficiencies; Evolution and Lifetimes. *ApJS* **181**, 321–350 (2009). 0811.1059.
  21. Crutcher, R. M., Hakobian, N. & Troland, T. H. Testing Magnetic Star Formation Theory. *ApJ* **692**, 844–855 (2009). 0807.2862.
  22. Bertram, E., Federrath, C., Banerjee, R. & Klessen, R. S. Statistical analysis of the mass-to-flux ratio in turbulent cores: effects of magnetic field reversals and dynamo amplification. *MNRAS* **420**, 3163–3173 (2012). 1111.5539.
  23. Zuckerman, B. & Evans, N. J., II. Models of massive molecular clouds. *ApJ* **192**, L149–L152 (1974).
  24. Larson, R. B. Turbulence and star formation in molecular clouds. *MNRAS* **194**, 809–826 (1981).
  25. Solomon, P. M., Rivolo, A. R., Barrett, J. & Yahil, A. Mass, luminosity, and line width relations of Galactic molecular clouds. *ApJ* **319**, 730–741 (1987).
  26. Elmegreen, B. G. & Scalo, J. Interstellar Turbulence I: Observations and Processes. *ARA&A* **42**, 211–273 (2004).
  27. Ballesteros-Paredes, J., Klessen, R. S., Mac Low, M.-M. & Vazquez-Semadeni, E. Molecular Cloud Turbulence and Star Formation. In Reipurth, B., Jewitt, D. & Keil, K. (eds.) *Protostars and Planets V*, 63–80 (2007).
  28. Dobbs, C. L. *et al.* Formation of Molecular Clouds and Global Conditions for Star Formation. *Protostars and Planets VI* 3–26 (2014). 1312.3223.
  29. Padoan, P. *et al.* The Star Formation Rate of Molecular Clouds. *Protostars and Planets VI* 77–100 (2014). 1312.5365.
  30. Crutcher, R. M., Hakobian, N. & Troland, T. H. Self-consistent analysis of OH Zeeman observations. *MNRAS* **402**, L64–L66 (2010). 0912.3024.
  31. Crutcher, R. M. Role of Magnetic Fields in Star Formation. *Highlights of Astronomy* **15**, 438–439 (2010).
  32. Fatuzzo, M. & Adams, F. C. Enhancement of Ambipolar Diffusion Rates through Field Fluctuations. In *American Astronomical Society Meeting Abstracts #200*, vol. 34 of *Bulletin of the American Astronomical Society*, 769 (2002).
  33. Heitsch, F., Zweibel, E. G., Slyz, A. D. & Devriendt, J. E. G. Magnetic Flux Transport in the ISM Through Turbulent Ambipolar Diffusion. *Ap&SS* **292**, 45–51 (2004).
  34. Kudoh, T., Basu, S., Ogata, Y. & Yabe, T. Three-dimensional simulations of molecular cloud fragmentation regulated by magnetic fields and ambipolar diffusion. *MNRAS* **380**, 499–505 (2007). 0706.2696.
  35. Kudoh, T. & Basu, S. Formation of Collapsing Cores in Subcritical Magnetic Clouds: Three-dimensional Magnetohydrodynamic Simulations with Ambipolar Diffusion. *ApJ* **728**, 123 (2011). 1012.5707.
  36. Vishniac, E. T. & Lazarian, A. Reconnection in the Interstellar Medium. *ApJ* **511**, 193–203 (1999).
  37. Eyink, G. *et al.* Flux-freezing breakdown in high-conductivity magnetohydrodynamic turbulence. *Nature* **497**, 466–469 (2013).
  38. McKee, C. F. Photoionization-regulated star formation and the structure of molecular



- clouds. *ApJ* **345**, 782–801 (1989).
39. Myers, P. C. & Goodman, A. A. Magnetic molecular clouds - Indirect evidence for magnetic support and ambipolar diffusion. *ApJ* **329**, 392–405 (1988).
  40. Bertoldi, F. & McKee, C. F. Pressure-confined clumps in magnetized molecular clouds. *ApJ* **395**, 140–157 (1992).
  41. Crutcher, R. M., Troland, T. H., Lazareff, B., Paubert, G. & Kazès, I. Detection of the CN Zeeman Effect in Molecular Clouds. *ApJ* **514**, L121–L124 (1999).
  42. Bourke, T. L., Myers, P. C., Robinson, G. & Hyland, A. R. New OH Zeeman Measurements of Magnetic Field Strengths in Molecular Clouds. *ApJ* **554**, 916–932 (2001). [astro-ph/0102469](#).
  43. Crutcher, R., Heiles, C. & Troland, T. Observations of Interstellar Magnetic Fields. In Falgarone, E. & Passot, T. (eds.) *Turbulence and Magnetic Fields in Astrophysics*, vol. 614 of *Lecture Notes in Physics*, Berlin Springer Verlag, 155–181 (2003).
  44. Troland, T. H. & Crutcher, R. M. Magnetic Fields in Dark Cloud Cores: Arecibo OH Zeeman Observations. *ApJ* **680**, 457–465 (2008). [0802.2253](#).
  45. Vázquez-Semadeni, E. *et al.* Molecular cloud evolution - IV. Magnetic fields, ambipolar diffusion and the star formation efficiency. *MNRAS* **414**, 2511–2527 (2011). [1101.3384](#).
  46. Körtgen, B. & Banerjee, R. Impact of magnetic fields on molecular cloud formation and evolution. *MNRAS* **451**, 3340–3353 (2015). [1502.03306](#).
  47. Heitsch, F. & Hartmann, L. Accretion and diffusion time-scales in sheets and filaments. *MNRAS* **443**, 230–240 (2014). [1406.2191](#).
  48. Passot, T., Vazquez-Semadeni, E. & Pouquet, A. A Turbulent Model for the Interstellar Medium. II. Magnetic Fields and Rotation. *ApJ* **455**, 536–+ (1995). [arXiv:astro-ph/9601182](#).
  49. Hartmann, L., Ballesteros-Paredes, J. & Bergin, E. A. Rapid Formation of Molecular Clouds and Stars in the Solar Neighborhood. *ApJ* **562**, 852–868 (2001). [arXiv:astro-ph/0108023](#).
  50. Fletcher, A., Beck, R., Shukurov, A., Berkhuijsen, E. M. & Horellou, C. Magnetic fields and spiral arms in the galaxy M51. *MNRAS* **412**, 2396–2416 (2011). [1001.5230](#).
  51. Planck Collaboration *et al.* Planck intermediate results. XXXV. Probing the role of the magnetic field in the formation of structure in molecular clouds. *ArXiv e-prints: 1502.04123* (2015). [1502.04123](#).
  52. Fryxell, B. *et al.* FLASH: An Adaptive Mesh Hydrodynamics Code for Modeling Astrophysical Thermonuclear Flashes. *ApJS* **131**, 273–334 (2000).
  53. Dubey, A. *et al.* Challenges of Extreme Computing using the FLASH code. In Pogorelov, N. V., Audit, E. & Zank, G. P. (eds.) *Numerical Modeling of Space Plasma Flows*, vol. 385 of *Astronomical Society of the Pacific Conference Series*, 145–+ (2008).
  54. Bouchut, F., Klingenberg, C. & Waagan, K. A multiwave approximate Riemann solver for ideal MHD based on relaxation. I: theoretical framework. *Numerische Mathematik* **108**, 7–42 (2007).
  55. Bouchut, F., Klingenberg, C. & Waagan, K. A multiwave approximate Riemann solver for ideal MHD based on relaxation II - Numerical implementation with 3 and 5 waves. *Numerische Mathematik* **115**, 647–679 (2010).

56. Waagan, K., Federrath, C. & Klingenberg, C. A robust numerical code for compressible magnetohydrodynamics and its application to highly supersonic turbulence. *J. Comp. Phys.*, *submitted* (2010).
57. Duffin, D. F. & Pudritz, R. E. Simulating hydromagnetic processes in star formation: introducing ambipolar diffusion into an adaptive mesh refinement code. *MNRAS* **391**, 1659–1673 (2008). 0810.0299.
58. Koyama, H. & Inutsuka, S.-i. An Origin of Supersonic Motions in Interstellar Clouds. *ApJ* **564**, L97–L100 (2002). arXiv:astro-ph/0112420.
59. Koyama, H. & Inutsuka, S.-I. Molecular Cloud Formation in Shock-compressed Layers. *ApJ* **532**, 980–993 (2000). arXiv:astro-ph/9912509.
60. Vázquez-Semadeni, E. *et al.* Molecular Cloud Evolution. II. From Cloud Formation to the Early Stages of Star Formation in Decaying Conditions. *ApJ* **657**, 870–883 (2007). astro-ph/0608375.
61. Banerjee, R., Vázquez-Semadeni, E., Hennebelle, P. & Klessen, R. S. Clump morphology and evolution in MHD simulations of molecular cloud formation. *MNRAS* **398**, 1082–1092 (2009). 0808.0986.
62. Federrath, C., Banerjee, R., Clark, P. C. & Klessen, R. S. Modeling Collapse and Accretion in Turbulent Gas Clouds: Implementation and Comparison of Sink Particles in AMR and SPH. *ApJ* **713**, 269–290 (2010). 1001.4456.
63. Truelove, J. K. *et al.* The Jeans Condition: A New Constraint on Spatial Resolution in Simulations of Isothermal Self-gravitational Hydrodynamics. *ApJ* **489**, L179+ (1997).
64. Vishniac, E. T. Nonlinear instabilities in shock-bounded slabs. *ApJ* **428**, 186–208 (1994). astro-ph/9306025.
65. Mouschovias, T. C. & Spitzer, L., Jr. Note on the collapse of magnetic interstellar clouds. *ApJ* **210**, 326–+ (1976).
66. Heitsch, F., Burkert, A., Hartmann, L. W., Slyz, A. D. & Devriendt, J. E. G. Formation of Structure in Molecular Clouds: A Case Study. *ApJ* **633**, L113–L116 (2005). arXiv:astro-ph/0507567.
67. Heitsch, F., Hartmann, L. & Burkert, A. Fragmentation of Shocked Flows: Gravity, Turbulence and Cooling. *ArXiv e-prints* **805** (2008). 0805.0801.
68. Inoue, T. & Inutsuka, S.-i. Two-Fluid Magnetohydrodynamic Simulations of Converging H I Flows in the Interstellar Medium. I. Methodology and Basic Results. *ApJ* **687**, 303–310 (2008). 0801.0486.
69. Field, G. B. Thermal Instability. *ApJ* **142**, 531–+ (1965).
70. Li, H.-B. *et al.* Evidence for dynamically important magnetic fields in molecular clouds. *MNRAS* **411**, 2067–2075 (2011). 1007.3312.
71. Padoan, P. *et al.* MHD Turbulence In Star-Forming Clouds. In Bertin, G., de Luca, F., Lodato, G., Pozzoli, R. & Romé, M. (eds.) *American Institute of Physics Conference Series*, vol. 1242 of *American Institute of Physics Conference Series*, 219–230 (2010).
72. Shu, F. H., Allen, A., Shang, H., Ostriker, E. C. & Li, Z.-Y. Low-Mass Star Formation: Theory. In Lada, C. J. & Kylafis, N. D. (eds.) *NATO Advanced Science Institutes (ASI) Series C*, vol. 540 of *NATO Advanced Science Institutes (ASI) Series C*, 193 (1999).
73. McKee, C. F., Zweibel, E. G., Goodman, A. A. & Heiles, C. Magnetic Fields in Star-Forming Regions - Theory. In Levy, E. H. & Lunine, J. I. (eds.) *Protostars and*

*Planets III*, 327–+ (1993).

74. Jülich Supercomputing Centre. JUQUEEN: IBM Blue Gene/Q Supercomputer System at the Jülich Supercomputing Centre. *Journal of large-scale research facilities* **1**, A1 (2015).